

High-Resolution Spatio-Temporal Precipitation Climatology in Complex Terrain

Reto Stauffer, Georg J. Mayr, Jakob Messner, Nikolaus Umlauf, Achim Zeileis
 Contact: reto.stauffer@uibk.ac.at

Introduction

High-resolution spatio-temporal climatologies of precipitation are important for e.g., agriculture, risk assessments, or tourism.

Problem: large proportion of **zero-observations** for high (daily) temporal resolutions (dry days). **A common solution:** decrease the temporal resolution to **monthly** or **yearly** sums to remove zeros (Goovaerts 2000, Hijmans 2005, ...).

Our goal

Provide a methodology to create climatologies with:

- **daily** temporal resolution (requires handling of zero-observations)
- **full climatological distribution**
- fully **scaleable** spatial domain/resolution
- **no need** for **extensive** or **manual** tuning for new areas

Model Setup

Generalized Additive Model for Location, Scale, and Shape

$$y^* \sim \mathcal{N}(\mu, \sigma^2), \quad y = \max(0, y^*), \quad \mu = \eta_\mu, \quad \log(\sigma) = \eta_\sigma$$

$$\eta = \beta_0 + \underbrace{\beta_1 \text{ alt}}_{\text{altitude}} + \underbrace{f_1(\text{yday})}_{\text{season}} + \underbrace{f_2(\text{long, lat})}_{\text{spatial}} + \underbrace{f_3(\text{yday, long, lat})}_{\text{spatial season}}$$

y^*/y :	latent/censored response	η :	linear predictor (identical for $\mu/\log(\sigma)$)
$\mathcal{N}(\dots)$:	Gaussian distribution	β :	unknown coefficients
$f_k(\dots)$:	non-linear multidimensional functions	yday:	day of the year
μ/σ :	mean and standard deviation	long/lat/alt:	geographical information

Precipitation (y) is physically limited to ≥ 0 , which can be considered by an e.g., censored distribution ($\max(0, y^*)$). A novel Bayesian model framework was used to estimate the unknown coefficients (*R* package **bamlss**, Umlauf 2015).

Data & Data Analysis

Data set

- 110 stations with quality controlled data; 510–2300m *a.m.s.l.*
- 24h sums observed at 06UTC
- 42 years of data (85% data availability)
- 1'440'000 observations; fraction of zeros $\sim 56\%$




Figure 1: Two sample stations; 0.90/0.75/0.50 quantiles are shown for both, the daily observed precipitation distribution over the last 33/24 years, and the in-sample fitted spatial climatology on a daily basis (final results).

Features captured by the statistical model

- seasonal pattern vary with location (e.g., north/south; Fig. 1)
- increased amount of precipitation during the summer season (Figs. 1&2)
- independent seasonal pattern for mean (μ) and variance (σ) (\updownarrow ; Figs. 1&2)
- significantly drier south of the main Alpine ridge (Figs. 1&3)

Acknowledgements:

Ongoing project funded by the **Austrian Science Fund (FWF)**: TRP 290-N26. The computational results presented have been achieved in part using the **Vienna Scientific Cluster (VSC)**. Data set provided by the "Ministerium für ein lebenswertes Österreich", **hydrographical service Tyrol** (ehyd.gv.at).

Model Results




Figure 2: Centered cyclic seasonal effect $f_1(\text{yday})$. "April weather" (blue): increasing mean (μ), local maximum of variability ($\log(\sigma)$). Autumn (orange): strongly decreasing mean (μ) with coincidentally increasing variability ($\log(\sigma)$).




Figure 3: Centered longitude/latitude effect $f_2(\text{long, lat})$. Mean (μ): positive effect north while dry to the South/inner-alpine regions. Variance ($\log(\sigma)$): weaker variability within the Alps.




Figure 4: Estimated expectation (*left*) and probability of precipitation (*right*), 3. September. **Innsbruck:** expected precipitation amount: 3.5mm, probability of precipitation: 46%. **Hafelekar:** expected precipitation amount: 6.4mm, probability of precipitation: 57%.

Summary & Outlook

Current status

- **zero-observations** properly handled by the **censored distribution**
- highly resolved **spatio-temporal** model; **adaptable** to arbitrary scales
- **full climatological distribution**; allows extracting **quantiles/probabilities**
- accurate estimate on **station level**
- "simple", **generalized** setup

Planned extensions

- to include additional **covariates** (e.g., terrain dependent features, wind)
- to test different **distributions** (e.g., censored logistic)
- to include **additional stations**
- to **compare** with existing methods

References:

Goovaerts, P., 2000: Geostatistical approaches for incorporating elevation into the spatial interpolation of rainfall. *Journal of Hydrology*, **228** (1–2), 113–129.
 Hijmans, R. J. et al., 2005: Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25** (15), 1965–1978.
 Stauffer, R., et al., 2015: Spatio-temporal Censored Model of Precipitation Climatology. *Proceedings of IWSM 2015*. Linz: Johannes Kepler University, Proceedings Book 2015 (1), 366–371.
 Umlauf, N., et al., 2015: bamlss: Bayesian Additive Models for Location Scale and Shape (and Beyond). *R package version 0.1-1* (https://R-Forge.R-project.org/R/?group_id=865).

